

## VACCINATIONS FOR ALL

- + **Flu Shots**
- + **Pediatric Pfizer Doses**
  - First and Second Doses
  - Additional Primary Series Doses for Those Who Are Immunocompromised
  - Booster Doses for People Ages 12 and Older
  - One-Third the Strength of the Adult Dose

### Available at any of our hospitals

With COVID-19 and flu season occurring together, it is more important than ever to protect yourself and your children from getting sick.

COVID-19 and flu vaccines are safe and highly effective in reducing risk of illness, hospitalization and even death.

Your children can get a COVID-19 vaccine and the flu shot at the same time with no out-of-pocket cost to you.

Learn more about which vaccine is right for you. Scan the QR code or call **1-844-NYC-4NYC** (1-844-692-4692).

Learn more  
about COVID-19  
Vaccines



Learn more  
about Flu  
Vaccine



# Who Should Get Vaccinated for Flu and COVID-19?

## **Influenza (Flu) Vaccine**

Everyone six months and older should get a seasonal influenza (flu) vaccine each year, especially:

- + Children under 2
- + People over 65
- + People who are pregnant or recently gave birth
- + People with medical conditions like diabetes, asthma, heart disease or HIV
- + Essential workers, including health care workers and caregivers of people that are at a higher risk of getting sick. Those who care for children, especially infants younger than six months, should also get the flu vaccine.

## **COVID-19 Vaccine**

Most individuals are now eligible to get a COVID-19 vaccine:

- + People aged 5 to 11 can get two pediatric doses (one-third strength) of Pfizer-BioNTech
- + People aged 12 and older can get two doses of the Pfizer-BioNTech vaccine

## **Additional Primary Series Doses of Pfizer COVID-19 Vaccine**

Additional primary series doses are available to people who got their second dose of COVID-19 vaccine at least 28 days ago and have a weakened immune system due to a condition or treatment. These may include:

- + Active treatment for cancer
- + Organ transplant and medicine to reduce the strength of the immune system
- + Stem cell transplant within the last 2 years
- + Moderate or severe primary immunodeficiency
- + Advanced or untreated HIV
- + Active treatment with a high dose of corticosteroids or other drugs that moderately or severely weaken the immune system

You must be 5 or older to get the Pfizer additional primary series dose. Additional primary series doses are not recommended for most people with other medical conditions such as diabetes, heart or lung disease.

## **COVID-19 Booster Dose - Pfizer and Moderna Vaccine Recipients**

Everyone 12 years or older should get a booster shot at least 5 months after receiving the second dose of their primary Pfizer COVID-19 vaccination series.

People ages 12-17 who previously got the Pfizer primary series more than 5 months ago are eligible for the Pfizer booster shot only.

People who are 12 or older and immunocompromised should get a booster shot at least 3 months after receiving the additional dose of their primary Pfizer COVID-19 vaccination series for a total of 4 doses.

Hospitals in the NYC Health + Hospitals system offer Pfizer and J&J vaccines. NYC Health + Hospitals/Gotham Health sites offer Moderna vaccine. For a full list of locations, visit our website: [nychealthandhospitals.org](https://nychealthandhospitals.org).