

VACCINATIONS FOR ALL

Pfizer and J&J COVID-19 Vaccines

- + Adult Doses
- + Doses for children 5 years and up
- + Additional Primary Series Doses for Those Who Are Immunocompromised
- + Boosters

Available here to all who are eligible

With COVID-19, it is more important than ever to protect yourself and loved ones from getting sick.

COVID-19 vaccines are safe and highly effective in reducing risk of illness, hospitalization and even death.

You can get a COVID-19 vaccine with no out-of-pocket cost to you.

Learn more about which vaccine is right for you. Scan the QR code or call **1-844-NYC-4NYC** (1-844-692-4692).

Learn more
about COVID-19
Vaccines



Learn more
about Flu
Vaccine



Who Should Get Vaccinated for COVID-19?



COVID-19 Vaccine

Most individuals are now eligible to get a COVID-19 vaccine:

- + People aged 5 to 11 can get two pediatric doses (one-third strength) of Pfizer-BioNTech
- + People 12 and older can get two doses of Pfizer-BioNTech
- + People 18 and older can get two doses of Moderna, or one dose of Johnson & Johnson/Janssen

Additional Primary Series Doses of Pfizer, Moderna or Johnson & Johnson COVID-19 Vaccine

Additional primary series doses of an mRNA vaccine are preferred for people who got their second dose of an mRNA COVID-19 vaccine or single dose of a Johnson & Johnson vaccine at least 28 days ago and have a weakened immune system due to a condition or treatment. These may include:

- + Active treatment for cancer
- + Organ transplant and medicine to reduce the strength of the immune system
- + Stem cell transplant within the last 2 years
- + Moderate or severe primary immunodeficiency
- + Advanced or untreated HIV
- + Active treatment with a high dose of corticosteroids or other drugs that moderately or severely weaken the immune system

You must be 5 or older to get the Pfizer additional primary series dose and 18 or older to get the Moderna additional primary series dose. In some situations, you can consider giving a Johnson & Johnson/Janssen vaccine as an additional primary series dose. Additional primary series doses are not recommended for most people with other medical conditions such as diabetes, heart or lung disease.

COVID-19 Booster Dose - Pfizer and Moderna Vaccine Recipients

Everyone 5 years or older should get a booster shot at least 5 months after receiving the second dose of their primary Pfizer or Moderna COVID-19 vaccination series.

People who received two shots of the Pfizer or Moderna vaccine should try to get the same vaccine as their booster, but have the choice to switch to any of the three types of vaccines authorized in the U.S.

People ages 5-17 who previously got the Pfizer primary series more than five months ago are eligible for the Pfizer booster shot only.

COVID-19 Booster Dose - Johnson & Johnson Vaccine Recipients

Everyone 18 years or older should get a booster shot at least two months after receiving the single dose primary COVID-19 vaccination. Pfizer or Moderna (mRNA COVID-19 vaccines) boosters are preferred in most situations.

Immunocompromised people who received J&J as a first dose and an mRNA vaccine at least 28 days later as an additional dose should get a booster dose at least 2 months after the additional dose of mRNA vaccine, for a total of 3 doses.

Hospitals in the NYC Health + Hospitals system offer Pfizer and J&J vaccines. NYC Health + Hospitals/Gotham Health sites offer Moderna vaccine and Pfizer vaccine for children 5-11 years of age. For a full list of locations, visit our website: nychealthandhospitals.org.