

VACCINATIONS FOR ALL

Pfizer COVID-19 Vaccines

- + Adult Doses
- + Doses for children 6 months and up
- + Additional Primary Series Doses for Those Who Are Immunocompromised
- + Booster shots

Available here to all who are eligible

With COVID-19, it is more important than ever to protect yourself and loved ones from getting sick.

COVID-19 vaccines are safe and highly effective in reducing risk of illness, hospitalization and even death.

You can get a COVID-19 vaccine with no out-of-pocket cost to you.

Learn more about which vaccine is right for you. Scan the QR code or call **1-844-NYC-4NYC** (1-844-692-4692)

Learn more
about COVID-19
Vaccines



Who Should Get Vaccinated for COVID-19?

- + Children ages 6 months to 4 years can get three pediatric doses (one-tenth strength) of Pfizer-BioNTech
- + Children ages 5 to 11 can get two pediatric doses (one-third strength) of Pfizer-BioNTech
- + Children and adults 12 and older can get two doses of Pfizer-BioNTech
- + Adults 18 and older can get two doses of Moderna, or one dose of Johnson & Johnson/Janssen
- + Children and adults 5 years and older who have a weakened immune system due to a condition or treatment can get an additional Pfizer primary series dose – also known as a third dose.

Who Should Get a COVID-19 Booster Shot?

Everyone 5 years or older should get a booster shot at least five months after receiving the second dose of their primary Pfizer or Moderna COVID-19 vaccination series.

People who received two shots of the Pfizer or Moderna vaccine should try to get the same vaccine as their booster, but they have the choice to switch to any of the three types of vaccines authorized in the U.S.

People ages 5 to 17 who previously got the Pfizer primary series more than five months ago are eligible for the Pfizer booster shot only.

Everyone 18 years or older who received the Johnson & Johnson vaccine should get a booster shot at least two months after receiving their primary dose. Pfizer or Moderna (mRNA COVID-19 vaccines) boosters are preferred in most situations.

Immunocompromised people who received J&J as a first dose and an mRNA vaccine at least 28 days later as an additional dose should get a booster dose of an mRNA vaccine at least two months after the additional dose of mRNA vaccine, for a total of three doses.

Hospitals in the NYC Health + Hospitals system offer Pfizer and J&J vaccines. NYC Health + Hospitals/Gotham Health sites offer Moderna vaccine and Pfizer vaccine for children 5 to 11 years of age. For a full list of locations, visit our website: nychealthandhospitals.org.